

Note to Readers: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Prenatal Phthalate, Perfluoroalkyl Acid, and Organochlorine Exposures and Term Birth Weight in Three Birth Cohorts: Multi- Pollutant Models Based on Elastic Net Regression

Virissa Lenters, Lützen Portengen, Anna Rignell-Hydbom, Bo A.G. Jönsson, Christian H. Lindh,
Aldert H. Piersma, Gunnar Toft, Jens Peter Bonde, Dick Heederik, Lars Rylander, and
Roel Vermeulen

Table of Contents

Figure S1. Directed acyclic graph to describe the authors' view of the relationships between the contaminant exposures, covariates and birth weight.

Table S1. Serum concentrations of exposure biomarkers in pregnant women from Greenland (n=513), Warsaw, Poland (n=180), and Kharkiv, Ukraine (n=557), 2002–2004.

Table S2. Spearman correlation coefficients between exposure biomarkers^a in pooled samples, and per study population.

Table S3. Adjusted associations (β [95% CI]) between term birth weight and selected exposures and demographic, reproductive, and lifestyle factors.

Table S4. Single-exposure unpenalized OLS-regression models for term birth weight (n=1250).

Table S5. Assessment of potential effect modification of the associations between contaminant exposures and term birth weight.

Figure S2. Generalized additive models for the single-pollutant exposure–outcome relationship for PFOS and term birth weight, fitted with a smoothing spline term for PFOS (with restricted maximum likelihood estimation).